Metaxalone 640mg Tablets

NDC 74157-0018-60 • 60ct

NEWLY FDA APPROVED

COPAY
$0

NON-controlled Muscle Relaxer

Migrate all your muscle relaxers to Metaxalone 640mg NDC 74157-0018-60

For Most Commercial Insured Patients*

*** Up to $100 off patient's copay per 30 day Rx or up to $300 per 90 day Rx**

An Estimated 126.6M Americans (1 In 2 Adults) Are Affected By Musculoskeletal Conditions That Include But Are Not Limited To:

• Back and/or neck pain • Carpal Tunnel Syndrome
• Osteoarthritis • Joint pain

INDICATIONS AND USAGE

Metaxalone Tablets, 640mg, are indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomforts associated with acute, painful musculoskeletal conditions in adults and pediatric patients 13 years of age and older.

DOSAGE AND ADMINISTRATION

The recommended dosage of Metaxalone Tablets, 640mg in adults and pediatric patients 13 years of age and older is 640mg taken orally with or without food, three to four times a day. The maximum recommended daily dosage is 2,560 mg (one tablet four times a day).

Metaxalone Tablets, 640mg and SKELAXIN 800mg are not mutually substitutable on a mg-to-mg basis due to differences in pharmacokinetic profiles [see Warnings and Precautions (5.3) and Clinical Pharmacology (12.3)]. When it is appropriate to switch a patient taking SKELAXIN 800mg on an empty stomach to Metaxalone Tablets, 640mg:

•Stop SKELAXIN 800mg three times a day and start Metaxalone Tablets 640mg three times a day on an empty stomach, OR
•Stop SKELAXIN 800mg four times a day and start Metaxalone Tablets 640mg four times a day on an empty stomach.

For patients who have been taking either product with food, do not switch between SKELAXIN 800mg and Metaxalone Tablets, 640mg.

ADVERSE REACTIONS

The following adverse reactions associated with the use of metaxalone were identified in clinical studies or postmarketing reports. Because some of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The most frequent reactions to metaxalone were:
•CNS: drowsiness, dizziness, headache, and nervousness or "irritability"
•Digestive: nausea, vomiting, gastrointestinal upset.

Other adverse reactions were:
•Allergic: anaphylaxis have been reported with metaxalone.
•CNS: cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of metaxalone (within the recommended dosage range) and other serotonergic drugs and with the use of metaxalone as the only serotonergic drug taken at a dosage higher than the recommended dosage [see Warnings and Precautions (5.1), Drug Interactions(7.2), and Overdosage (10)].
•Hematologic: leucopenia; hemolytic anemia.
•Hepatobiliary: jaundice.
•Immune System: hypersensitivity reaction, rash with or without pruritus.

Manufacture direct pricing

Description	NDC	AWP	Package Size
Metaxalone 640mg	74157-0018-60	$4,500.0	60 ct